**Product Sheet** 

# Endothelial Cell Growth Kit-VEGF

PCS-100-041<sup>™</sup>

### Description

The Endothelial Cell Growth Kit-VEGF contains components that when added to Vascular Cell Basal Medium (ATCC PCS-100-030) create a complete, low serum culture environment for human large vessel endothelial cells. Use of the Endothelial Cell Growth Kit-VEGF will support a faster rate of proliferation because of the presence of several purified human recombinant (rh) growth factors (rh VEGF, rh EGF, rh FGF basic, and rh IGF-1) combined with heparin and hydrocortisone. **Shipping information:** 1 kit

### **Storage Conditions**

Product format: Frozen Storage conditions: -20°C or colder, -70°C for long-term storage

### Intended Use

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use.

## BSL 1

ATCC determines the biosafety level of a material based on our risk assessment as guided by the current edition of *Biosafety in Microbiological and Biomedical Laboratories (BMBL)*, U.S. Department of Health and Human Services. It is your responsibility to understand the hazards associated with the material per your organization's policies

PCS-100-041

and procedures as well as any other applicable regulations as enforced by your local or national agencies.

### **Certificate of Analysis**

For batch-specific test results, refer to the applicable certificate of analysis that can be found at www.atcc.org.

### Handling Procedures

#### **Unpacking and Storage Instructions**

- 1. Check all containers for leakage or breakage.
- 2. Store the basal medium at 2°C to 8°C and the growth kit(s) at either -20°C in a freezer that is not self-defrosting or at -70°C for long-term storage. If thawed upon arrival, the growth kit should be stored at 2°C to 8°C and added to the basal medium within 72 hours of receipt.

#### **Preparation of Complete Growth Media**

- 1. Obtain one growth kit from the freezer; make sure that the caps of all components are tight.
- 2. Thaw the components of the growth kit just prior to adding them to the basal medium. It is necessary to warm the L-glutamine component in a 37°C water bath and shake to dissolve any precipitates, prior to adding to the basal medium.
- 3. Obtain one bottle of Vascular Cell Basal Medium (475 mL) from cold storage.
- 4. Decontaminate the external surfaces of all growth kit component vials and the basal medium bottle by spraying them with 70% ethanol.
- Using aseptic technique, and working in a laminar flow hood or biosafety cabinet transfer <u>the volume of</u> each growth kit component, as indicated in Table 1 or 2, to the bottle of basal medium using a separate sterile pipette for

PCS-100-041

each transfer.

6. Tightly cap the bottle of complete growth medium and swirl the contents gently to assure a homogeneous solution. Do not shake forcefully to avoid foaming. Label and date the bottle.

Complete media should be stored in the dark at 2°C to 8°C (do not freeze). When stored under these conditions, complete media is stable for 30 days.

Table 1. If using	the Endothelial	Cell Growth	Kit-BBE (ATCC <sup>®</sup>	PCS-100-040),	add the
indicated volume	for each compo	onent:			

Component	Volume	Final Concentration
Bovine Brain Extract (BBE)	1.0 mL	0.2%
rh EGF	0.5 mL	5 ng/mL
L-glutamine	25.0 mL	10 mM
Heparin sulfate	0.5 mL	0.75 Units/mL
Hydrocortisone hemisuccinate	0.5 mL	1 µg/mL
Fetal Bovine Serum	10.0 mL	2%
Ascorbic acid	0.5 mL	50 μg/mL

Table 2. If using the Endothelial Cell Growth Kit-VEGF (ATCC $^{\otimes}$  PCS-100-041), add the indicated volume for each component:

Component	Volume	Final Concentration
-----------	--------	------------------------



PCS-100-041

rh VEGF	0.5 mL	5 ng/mL
rh EGF	0.5 mL	5 ng/mL
rh FGF basic	0.5 mL	5 ng/mL
rh IGF-1	0.5 mL	15 ng/mL
L-glutamine	25.0 mL	10 mM
Heparin sulfate	0.5 mL	0.75 Units/mL
Hydrocortisone hemisuccinate	0.5 mL	1 μg/mL
Fetal Bovine Serum	10.0 mL	2%
Ascorbic acid	0.5 mL	50 μg/mL

Antimicrobials and phenol red are not required for proliferation but may be added if desired. The recommended volume of each optional component to be added to the complete growth media is summarized in Table 3.

Table 3. Addition of Antimicrobials/Antimycotics and Phenol Red (Optional)

Component	Volume	Final Concentration
Gentamicin- Amphotericin	0.5 mL	Gentamicin: 10 µg/mL
B Solution		Amphotericin Β: 0.25 μg/mL



PCS-100-041

Penicillin- Streptomycin-	0.5 mL	Penicillin: 10 Units/mL
Amphotericin B Solution		Streptomycin: 10 µg/mL
		Amphotericin B: 25 ng/mL
Phenol Red	0.5 mL	33 µM

### **Quality Control Specifications**

Bacterial and fungal testing: Not detected Mycoplasma contamination: Not detected Functional tests: Rate of proliferation and morphology

### **Material Citation**

If use of this material results in a scientific publication, please cite the material in the following manner: Endothelial Cell Growth Kit-VEGF (ATCC PCS-100-041)

#### References

References and other information relating to this material are available at www.atcc.org.

### Warranty

The product is provided 'AS IS' and the viability of  $\text{ATCC}^{\otimes}$  products is warranted for 30

**Product Sheet** 

PCS-100-041

days from the date of shipment, provided that the customer has stored and handled the product according to the information included on the product information sheet, website, and Certificate of Analysis. For living cultures, ATCC lists the media formulation and reagents that have been found to be effective for the product. While other unspecified media and reagents may also produce satisfactory results, a change in the ATCC and/or depositor-recommended protocols may affect the recovery, growth, and/or function of the product. If an alternative medium formulation or reagent is used, the ATCC warranty for viability is no longer valid. Except as expressly set forth herein, no other warranties of any kind are provided, express or implied, including, but not limited to, any implied warranties of merchantability, fitness for a particular purpose, manufacture according to cGMP standards, typicality, safety, accuracy, and/or noninfringement.

### Disclaimers

This product is intended for laboratory research use only. It is not intended for any animal or human therapeutic use, any human or animal consumption, or any diagnostic use. Any proposed commercial use is prohibited without a license from ATCC.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate or complete and the customer bears the sole responsibility of confirming the accuracy and completeness of any such information.

This product is sent on the condition that the customer is responsible for and assumes all risk and responsibility in connection with the receipt, handling, storage, disposal, and use of the ATCC product including without limitation taking all appropriate safety and handling precautions to minimize health or environmental risk. As a condition of receiving the material, the customer agrees that any activity undertaken with the ATCC product and any progeny or modifications will be conducted in compliance with all applicable laws, regulations, and guidelines. This product is provided 'AS IS' with no representations or warranties whatsoever except



www.atcc.org

Page 6 of 7

PCS-100-041

as expressly set forth herein and in no event shall ATCC, its parents, subsidiaries, directors, officers, agents, employees, assigns, successors, and affiliates be liable for indirect, special, incidental, or consequential damages of any kind in connection with or arising out of the customer's use of the product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the material transfer agreement (MTA) for further details regarding the use of this product. The MTA is available at www.atcc.org.

#### **Copyright and Trademark Information**

© ATCC 2023. All rights reserved.

ATCC is a registered trademark of the American Type Culture Collection.

### Revision

This information on this document was last updated on 2022-11-05

### **Contact Information**

ATCC 10801 University Boulevard Manassas, VA 20110-2209 USA US telephone: 800-638-6597 Worldwide telephone: +1-703-365-2700 Email: tech@atcc.org or contact your local distributor

