

## CELL DEATH PATHWAY BCL-2 FAMILY CELL LINE PANELS 1 & 2

The Cell Death Pathway BCL-2 Family Cell Line Panels 1 & 2 are useful for the study of cell apoptosis signaling pathways, BCL-2 family member molecular mechanisms, and mitochondrial dysfunction. Panel 1 (ATCC® <u>TCP-2100</u>™) comprises five cell lines that stably overexpress either wild-type BCL-2 or a mutant form of BCL-2. Panel 2 (ATCC® <u>TCP-2110</u>™) is composed of seven immortalized mouse embryonic fibroblast cell lines generated from a set of BCL-2 family member gene knockout mice. The table below provides more information for the cell lines included in these panels.

	ATCC® No.	Designation	Source	Cell type	Transfected gene	Significant features	Verification
BCL-2 Family Cell Panel 1 (ATCC® TCP-2100™)	<u>CRL-2898</u> ™	Neo Jurkat	acute T cell leukemia	Immortalized T lymphocyte	Empty vector	Normal expression of wild-type BCL-2	ATCC in-house sequencing
	<u>CRL-2899</u> ™	BCL2 Jurkat	acute T cell leukemia	Immortalized T lymphocyte	BCL-2	Stable overexpression of anti-apoptosis gene BCL-2	ATCC in-house sequencing
	<u>CRL-2900</u> ™	BCL2 (S70A) Jurkat	acute T cell leukemia	Immortalized T lymphocyte	BCL2 (S70A)	Stable overexpression of BCL-2 with a mutation at phosphory-lation site Ser70	ATCC in-house sequencing
	<u>CRL-2901</u> ™	BCL2 (S87A) Jurkat	acute T cell leukemia	Immortalized T lymphocyte	BCL2 (S87A)	Stable overexpression of BCL-2 with a substitution mutation at phosphorylation site Ser87	ATCC in-house sequencing
	<u>CRL-2902</u> ™	BCL2 (AAA) Jurkat	acute T cell leukemia	Immortalized T lymphocyte	BCL2 (AAA)	Stable overexpression of BCL-2 with a triple substitution at phosphorylation site Thr69, Ser70, and Ser87	ATCC in-house sequencing

	ATCC® No.	Designation	Source	Cell type	Genotype	Significant features	Verification
BCL-2 Family Cell Panel 2 (ATCC <sup>®</sup> <u>TCP-2110</u> ")	<u>CRL-2907</u> ™	WT SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	Wild type	Immortalized MEFs.	ATCC in-house qPCR test
	<u>CRL-2908</u> ™	BCL2 KO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BCL-2 knockout	Immortalized MEFs generated from anti- apoptosis BCL-2 gene knockout mice.	ATCC in-house qPCR test
	<u>CRL-2909</u> ™	BAD KO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BAD knockout	Immortalized MEFs generated from pro-apoptosis BAD gene knockout mice.	ATCC in-house qPCR test
	<u>CRL-2910</u> ™	BAX KO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BAX knockout	Immortalized MEFs generated from pro-apoptosis BAX gene knockout mice.	ATCC in-house qPCR test
	<u>CRL-2911</u> ™	BID KO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BID knockout	Immortalized MEFs generated from pro-apoptosis BID gene knockout mice.	ATCC in-house qPCR test
	<u>CRL-2912</u> ™	BAK KO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BAK knockout	Immortalized MEFs generated from pro-apoptosis BAK gene knockout mice	ATCC in-house qPCR test
	<u>CRL-2913</u> ™	BAX BAK DKO SV40 MEF	Embryo	immortalized mouse embryonic fibroblast	BAX & BAK double knockout	Immortalized MEFs generated from pro-apoptosis BAX and BAK double knockout mice. The cells are resistant to multiple apoptotic stimuli.	ATCC in-house qPCR test











## CB-122021-v02

©2022 American Type Culture Collection. The ATCC trademark and trade name, and any other trademarks listed in this publication are trademarks owned by the American Type Culture Collection unless indicated otherwise.