



## PARASITIC PROTOZOA GENOMIC DNA

Standardized molecular reagents are critical in the development of molecular assays used to diagnose parasitic disease. Genomic DNA preparations from parasitic protozoa provide researchers with rapid access to protozoan nucleic acids without the hassle of *in vitro* expansions. They are useful in a variety of applications, including PCR-based assays and whole genome sequencing.

ATCC offers a growing assortment of DNA prepared from our vast collection of living stock protists. Protozoa DNA are available in two

easy formats, including the ATCC Protozoa DNA Panels as well as individual reagents. Each preparation provides  $\geq 2 \mu\text{g}$  genomic DNA, tested for:

- Purity using absorbance
- Concentration by absorbance or PicoGreen®
- Integrity via gel electrophoresis
- Identity and functional activity by PCR amplification of the 18S rRNA gene or other gene targets

### ATCC Protozoa DNA Panels

The ATCC Protozoa DNA Panels are comprised of nucleic acid preparations isolated from taxonomic and medically-relevant strains. Each panel is offered at bulk discount pricing, allowing you to take advantage of added cost savings when compared to individual items.

#### LEISHMANIA DNA PANEL (ATCC® MP-13™)

Leishmaniasis is caused by parasites of the genus *Leishmania* that are transmitted via the bite of various phlebotomine sandfly species. Vaccines against leishmaniasis are still under development, thus the control of this disease relies on prompt diagnosis and chemotherapy in infected humans. The ATCC *Leishmania* DNA Panel (ATCC® MP-13™) consists of:

ATCC® No.	Description	Strain
50134D™	Genomic DNA from <i>Leishmania infantum</i>	MHOM/TN/80/IPT-1
PRA-309D™	Genomic DNA from <i>Leishmania major</i>	Seidman
30012D™	Genomic DNA from <i>Leishmania major</i>	—
30030D™	Genomic DNA from <i>Leishmania donovani</i>	Khartoum

#### ENTERIC PROTOZOA DNA PANEL (ATCC® MP-14™)

Diarrheal diseases due to intestinal protozoan pathogens cause a significant amount of morbidity and mortality worldwide. Three pathogens, *Giardia intestinalis*, *Cryptosporidium* sp., and *Entamoeba histolytica*, are responsible for the majority of intestinal disease caused by this group of pathogens. The ATCC Enteric Protozoa DNA Panel (ATCC® MP-14™) consists of:

ATCC® No.	Description	Strain
50803D™	Genomic DNA from <i>Giardia intestinalis</i>	WB Clone C6
PRA-67DQ™	Genomic DNA from <i>Cryptosporidium parvum</i>	Iowa
50608D™	Genomic DNA from <i>Blastocystis hominis</i>	BT1
30459DQ™	Genomic DNA from <i>Entamoeba histolytica</i>	HM-1:IMSS*

\*This strain only yields  $\geq 50$  ng of DNA.

# Individual Reagents

Select protozoa DNA preparations from our expanding list of selections, including DNA from *Neospora*, *Toxoplasma*, and *Trypanosoma*.

Species and ATCC® No.	Description	Strain
<b>ACANTHAMOEBA SP.</b>		
<a href="#">30010D™</a>	Genomic DNA from <i>Acanthamoeba castellanii</i>	Neff
<b>BLASTOCYSTIS HOMINIS</b>		
<a href="#">50177D™</a>	Genomic DNA from <i>Blastocystis hominis</i>	Nand II
<a href="#">50608D™</a>	Genomic DNA from <i>Blastocystis hominis</i>	BT1
<b>CRYPTOSPORIDIUM SP.</b>		
<a href="#">PRA-67D™</a>	Genomic DNA from <i>Cryptosporidium parvum</i>	Iowa
<b>ENTAMOEBA HISTOLYTICA</b>		
<a href="#">30459D™</a>	Genomic DNA from <i>Entamoeba histolytica</i>	HM-1:IMSS*
<b>GIARDIA INTESTINALIS</b>		
<a href="#">30888D™</a>	Genomic DNA from <i>Giardia intestinalis</i>	Portland-1
<a href="#">30957D™</a>	Genomic DNA from <i>Giardia intestinalis</i>	WB
<a href="#">50803D™</a>	Genomic DNA from <i>Giardia intestinalis</i>	WB Clone C6
<b>LEISHMANIA SP.</b>		
<a href="#">50134D™</a>	Genomic DNA from <i>Leishmania infantum</i>	MHOM/TN/80/IPT-1
<a href="#">PRA-309D™</a>	Genomic DNA from <i>Leishmania major</i>	Seidman
<a href="#">30012D™</a>	Genomic DNA from <i>Leishmania major</i>	--
<a href="#">30030D™</a>	Genomic DNA from <i>Leishmania donovani</i>	Khartoum
<b>NEOSPOA SP.</b>		
<a href="#">50843D™</a>	Genomic DNA from <i>Neospora caninum</i>	Nc-1
<b>TOXOPLASMA GONDII</b>		
<a href="#">50174D™</a>	Genomic DNA from <i>Toxoplasma gondii</i>	RH
<b>TRYPANOSOMA SP.</b>		
<a href="#">30022D™</a>	Genomic DNA from <i>Trypanosoma lewisi</i>	Lincicome
<a href="#">30266D™</a>	Genomic DNA from <i>Trypanosoma cruzi</i>	Tulahuen
<a href="#">50823D™</a>	Genomic DNA from <i>Trypanosoma cruzi</i>	SYLVIO-X10
<b>TRICHOMONAS SP.</b>		
<a href="#">30001D™</a>	Genomic DNA from <i>Trichomonas vaginalis</i>	C-1:NIH

\*This strain only yields ≥50 ng of DNA.

The ATCC Protozoa & Algae webpage has a new look!

VISIT US ONLINE AT [ATCC.ORG/PRODUCTS/CELLS\\_AND\\_MICROORGANISMS/PROTOZOA.ASPX](https://www.atcc.org/products/cells_and_microorganisms/protozoa.aspx) TO LEARN MORE ABOUT:

- ATCC living stock protists classified by supergroup
- Protozoan nucleic acids
- Culture media

## PHONE

800.638.6597  
703.365.2700

## EMAIL

tech@atcc.org

## WEB

www.atcc.org



10801 University Blvd.  
Manassas, VA 20110

MB-078-03

© 2018 American Type Culture Collection. The ATCC trademark and trade name, any and all ATCC catalog numbers and any other trademarks listed in this publication are trademarks of the American Type Culture Collection unless indicated otherwise. The Polymerase Chain Reaction (PCR) process is covered by patents owned by Hoffman-La Roche, Inc. Use of the PCR process requires a license.

These products are for laboratory use only. Not for human or diagnostic use. ATCC products may not be resold, modified for resale, used to provide commercial services or to manufacture commercial products without prior ATCC written approval.