ATCC

Product Sheet

# Keratinocyte Growth Kit (ATCC® PCS-200-040™)

**Please read this FIRST**

Storage Temp.
**-20°C (or -70°C for long-term storage)**

Biosafety Level
**1**

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

*Or contact your local distributor*

## Description

**Product Description:** Keratinocyte Growth Kit (ATCC® PCS-200-040) contains components that when added to Dermal Cell Basal Medium (ATCC® PCS-200-030) create a complete ATCC® Primary Cell Solutions™ culture environment for keratinocytes derived from normal human epidermis (e.g., Primary Epidermal Keratinocytes, Normal, Human Neonatal Foreskin, ATCC® PCS-200-010). The complete serum-free (although not animalfree) media is formulated to inhibit fibroblast growth, and the low calcium concentration (0.06 mM) slows differentiation. No feeder layers, extracellular matrix proteins or other substrates are required. The only animal-derived component is bovine pituitary extract (BPE); if necessary, keratinocytes can be cultured in the absence of BPE.
**Volume:** 1 Kit

## Directions for Use

**Unpacking and Storage Instructions**

1. Check all containers for leakage or breakage.
2. Store the growth kit(s) at either -20°C in a freezer that is not self-defrosting or at -70°C for long-term storage. If thawed upon arrival, the Growth Kit can be stored at 2°C to 8°C and added to the basal medium within 72 hours of receipt.

**Preparation of Complete Growth Media**

1. Obtain one growth kit from the freezer; make sure that the caps of all components are tight.
2. Thaw the components of the growth kit just prior to adding to the basal medium. It is necessary to warm the L-glutamine component in a 37°C water bath and shake to dissolve any precipitates, prior to adding to the basal medium.
3. Obtain one bottle of Dermal Cell Basal Medium (485 mL) from cold storage.
4. Decontaminate the external surfaces of all growth kit component vials and the basal medium bottle by spraying with 70% ethanol.
5. Using aseptic technique and working in a laminar flow hood or biosafety cabinet, transfer the indicated volume of each growth kit component to the bottle of basal medium using a separate sterile pipette for each.
6. Tightly cap the bottle of complete growth medium and swirl the contents gently to assure a homogeneous solution. Do not shake forcefully to avoid foaming. Label and date the bottle.

Complete media should be stored in the dark at 2°C to 8°C (do not freeze). When stored under these conditions, complete media is stable for 30 days.

**Table 1.** Keratinocyte Growth Kit Components

| Component | Volume | Final Concentration |
|---|---|---|
| Bovine Pituitary Extract (BPE) | 2.0 mL | 0.4% |
| rh TGF-a | 0.5 mL | 0.5 ng/mL |
| L-Glutamine | 15.0 mL | 6 mM |
| Hydrocortisone Hemisuccinate | 0.5 mL | 100 ng/mL |
| rh Insulin | 0.5 mL | 5 µg/mL |
| Epinephrine | 0.5 mL | 1.0 µM |
| Apo-Transferrin | 0.5 mL | 5 µg/mL |

Antimicrobials and phenol red are not required for proliferation, but may be added if desired. The recommended volume of each ***optional*** component to be added to the complete growth media is summarized in Table 2.

**Table 2.** Addition of Antimicrobials/Antimycotics and Phenol Red (Optional)

| Component | Volume | Final Concentration |
|---|---|---|
| Gentamicin-Amphotericin B Solution | 0.5 mL | Gentamicin: 10 µg/mL<br>Amphotericin B: 0.25 µg/mL |
| Penicillin-Streptomycin-Amphotericin B Solution | 0.5 mL | Penicillin: 10 Units/mL<br>Streptomycin: 10 µg/mL<br>Amphotericin B: 25 ng/mL |
| Phenol Red | 0.5 mL | 33 µM |

## ATCC Warranty

Product Sheet

# Keratinocyte Growth Kit (ATCC® PCS-200-040™)

**Please read this FIRST**

Storage Temp.
**-20°C (or -70°C for long-term storage)**

Biosafety Level
**1**

The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

## Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.

© ATCC 2014. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [04/15]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

*Or contact your local distributor*