ATCC® | Credible leads to Incredible™

Technical Data Sheet: PC-3-Luc2

ATCC® Number	CRL-1435-LUC2™
Organism	*Homo sapiens*
Tissue/Disease Source	Prostate adenocarcimoma
Product Description	This luciferase expressing cell line was derived from PC-3 cell line by transduction with lentiviral vector encoding firefly luciferase gene (luc2) and subsequently through single cell cloning. • Signal noise ratio: ≥ 1,000 • Bioluminescence: ≥ 100,000 photons/cell/sec (subject to imaging and culture condition) • Confirmed to be murine pathogen free
Application	Excellent signal/background ratio and stable Luciferase expression make this cell line ideal for in vivo bioluminescence imaging of xenograft animal model to study human cancer and monitor activity of anti-cancer drug. It also can be used in cell-based assays for cancer research.

In vivo Bioluminescent Imaging



Figure 1: *In vivo* detection of luciferase activity of PC-3-Luc2. PC-3-Luc2 cells ($3x10^6$) were injected subcutaneously into the dorsal region near the thigh of male NSG mice. Tumor growth was monitored weekly using a Xenogen IVIS Spectrum. *In vivo* bioluminescence imaging demonstrated the progression of tumors.

Cell Morphology



Figure 2: Cell morphology of PC-3 parental and PC-3-Luc2. Cells were maintained in ATCC recommended culture conditions. Cell morphology was observed under microscopy and images were captured by digital camera.

Luciferase Expression



Figure 3: Linearity of luminescence and of *in vitro* quantification of luciferase activity of PC-3-Luc2. Cells were seeded in a 96-well plate at indicated cell numbers per well, and Bright-Glo (Promega) was added to the indicated wells. The luminescence of the plate was read within 10 minutes using a luminescence plate reader (A) and determined to have a linear correlation of bioluminescence intensity with cell numbers. (B) The plate was imaged using a Xenogen IVIS Spectrum to quantify that photons emitted per cell.