Product Sheet

Campylobacter sputorum biovar sputorum (ATCC® 49916™)

Please read this FIRST

Storage Temp.
2°C to 8°C

Biosafety Level
2

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Campylobacter sputorum biovar sputorum* (ATCC® 49916™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Designation: LRA 116.06.89
Deposited Name: *Campylobacter sputorum* subsp. *bubulus* (Florent) Veron and Chatelain

Propagation

Medium
ATCC® Medium 18: Trypticase Soy Agar/Broth
ATCC® Medium 260: Trypticase soy agar/broth with defibrinated sheep blood

Growth Conditions
Temperature: 37°C
Atmosphere: Microaerophilic

Propagation Procedure
1. Open vial according to enclosed instructions or visit www.atcc.org for instructions.
2. Rehydrate the entire pellet with approximately 0.5 mL of #18 broth.
3. Aseptically transfer the entire contents to a 5-6 mL tube of #18 broth. Additional test tubes can be inoculated by transferring 0.5 mL of the primary broth tube to these secondary broth tubes.
4. Use several drops of the primary broth tube to inoculate a #260 plate and/or #260 agar slant.
5. Or, to obtain a biphasic culture, add several drops of the primary broth tube to a #260 agar slant. Best practice is to incubate these slants at an angle.
6. Incubate at 37°C under microaerophilic conditions for 1-3 days. Use an anaerobe jar with an active catalyst and a microaerophilic gas generator pack or other acceptable method. All tubes and slants should be incubated with caps loosened.

Notes

This organism requires moist conditions for best growth. Growth at the broth/agar interface of the biphasic slant should occur within 1-2 days, but little turbidity will be seen. To observe growth, examine a wet mount of the broth under phase microscopy.
The cells do not Gram stain well using traditional procedures. To obtain the best results, use a basic fuchsin counterstain in place of the safranin.
Once good growth is obtained, transfer or freeze the culture. Adding an equal amount of 20% sterile glycerol to pooled broth from several biphasic slants, followed by freezing in liquid nitrogen or "ultra-low temperature" freezer is recommended.
Additional information on this culture is available on the ATCC web site at www.atcc.org.

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans.

Product Sheet

Campylobacter sputorum biovar sputorum (ATCC® 49916™)

Please read this FIRST



Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Campylobacter sputorum biovar sputorum* (ATCC® 49916™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor