



Product Sheet

Fusarium falciforme (ATCC® MYA-3636™)

Please read this **FIRST**



Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section



Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Fusarium falciforme* (ATCC® MYA-3636™)

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor

Description

Strain Designation: FS1

Product Description: An ampoule containing viable cells (yeast cells, spores, or agar cubes with mycelia) suspended in cryoprotectant.

Propagation

The information recommended in this section is to assist users in obtaining living culture(s) for their studies. The recommendation does not imply that the conditions or procedures provided below are optimum. Experienced researchers may initiate the growth of a culture in their own way.

ATCC® Medium 336: Potato dextrose agar (PDA)

ATCC® Medium 307: Cornmeal agar

ATCC® Medium 28: Emmons' modification of Sabouraud's agar

Growth Conditions

Temperature: 25°C to 30°C

Atmosphere: Typical aerobic

Recommended Procedure

Frozen ampoules packed in dry ice should either be thawed immediately or stored in liquid nitrogen. If liquid nitrogen storage facilities are not available, frozen ampoules may be stored at or below -70°C for approximately one week. **Do not under any circumstance store frozen ampoules at refrigerator freezer temperatures (generally -20°C).** Storage of frozen material at this temperature will result in the death of the culture.

1. To thaw a frozen ampoule, place in a **25°C to 30°C** water bath, until just thawed (**approximately 5 minutes**). Immerse the ampoule just sufficient to cover the frozen material. Do not agitate the ampoule.
2. Immediately after thawing, wipe down ampoule with 70% ethanol and aseptically transfer at least 50 µL (or 2-3 agar cubes) of the content onto a plate or broth with medium recommended.
3. Incubate the inoculum/strain at the temperature and conditions recommended. Inspect for growth of the inoculum/strain regularly. The sign of viability is noticeable typically after 2-4 days of incubation. However, the time necessary for significant growth will vary from strain to strain.

Colony and Cell Morphology: Colonies growing rapidly forming white aerial mycelium, green to bluish-brown when sporodochia are present; reverse usually colorless. Monophialides with a rather distinct collarette. Macroconidia usually moderately curved, with short, blunt apical and indistinctly pedicellate basal cells, mostly 3-septate, 28-42 × 4-6 µm. Microconidia usually abundant, 8-16 × 2.0-4.5 µm. Chlamydo spores common.

Notes

No special notes

Additional, updated information on this product may be available on the ATCC® web site at www.atcc.org.

DNA Sequence

18S ribosomal RNA gene, partial sequence; internal transcribed spacer 1, 5.8S ribosomal RNA gene, and internal transcribed spacer 2, complete sequence; and 28S ribosomal RNA gene, partial sequence
GGTCTCCGTTGGTGAACCGAGCGGAGGGATCATTACCGAGTTATACAACCTCATCAACCCTGTGAACATA
CCTATAACGTTGCCTCGGCGGGAACAGACGGCCCCGTAACACGGGCGCCCCGCCAGAGGACCCCC
TAACCTGTGTTCTATAATGTTTCTTCTGAGTAAACAAGCAAATAAAATTAACCTTTCAACAACGGATCTC
TTGGCTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAATGTGAATTCAGAATTCAGTGA
ATCATCGAATCTTTGAACGCACATTGCGCCCCGAGTATTCTGGCGGGCATGCCTGTTTCGAGCGTCATT
ACAACCCTCAGGCCCGGGCTGGCGTTGGGGATCGGCGGAAGCCCCCTGCGGGCACAACCGCGTC
CCCCAAATACAGTGGCGGTCCCGCCGAGCTTCCATTGCGTAGTAGCTAACACCTCGCAACTGGAGAG
CGGCGCGGCCACGCCGTA AACACCCA ACTTCTGAATGTTGACCTCGAATCAGGTAGGAATACCCGCT
GAACCTAAGCATATCAATAA

D1D2 region of the 28S ribosomal RNA gene

ATATCAATAAGCGGAGGAAAAGAAACCAACAGGGATTGCCCCAGTAACGGCGAGTGAAGCGGCAAC
AGCTCAAATTTGAAATCTGGCTCTCGGGCCCGAGTTGTAATTTGTAGAGGATGCTTTTGGTGAGGTGCC
TTCCGAGTTCCTGGAACGGGACGCCATAGAGGGTGAGAGCCCCGCTGGTTGGACACCGATCCTCTG



Product Sheet

Fusarium falciforme **(ATCC® MYA-3636™)**

Please read this **FIRST**

Storage Temp.
Frozen: -80°C or colder
Freeze-Dried: 2°C to 8°C
Live Culture: See Propagation Section

 Biosafety Level
1

Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: *Fusarium falciforme* (ATCC® MYA-3636™)

TAAAGCTCCTTCGACGAGTCGAGTAGTTTGGGAATGCTGCTCTAAATGGGAGGTATATGTCTTCTAAAG
CTAAATACCGGCCAGAGACCGATAGCGCACAAAGTAGAGTGATCGAAAGATGAAAAGAACCTTTGAAA
AGAGAGTTAAAAAGTACGTGAAATTTGTTGAAAGGGAAGCGCTTGTGACCAGACTTGGGCTTGGTTGA
TCATCCGGGGTTCTCCCCGGTGCCTCTCCGGCTCAGGCCAGCATCAGTTCGCCCTGGGGGATAAAGG
CTTCGGGAATGTGGCTCTCTCCGGGAGTGTATAGCCCGCTGCGTAATACCCTGTGGCGGACTGAGGT
TCGCGCATTGCAAGGATGCTGGCGTAATGGTCATCAGTGAC

Elongation factor 1-alpha

TTGGTGACATCTCCCCGATCGCGCCTTGCTATTCCACATCGAATTCCTCCCTCGCGATACGCTCTGCG
CCCCCTTCTCCCGAGTCCCAAAATTTTTCGGTCCGACCGTAATTTTGGTGGGGCATTACCCCGCC
ACTCGGGCGACTTGGACAAAGCCCTGATCCCTGCACACAAAAACACCAAACCTCTTGGCGCGCAT
CATCACGTGGTTTACAACAGACGCTAACCGGTCCAACAATAGGAAGCCGCTGAGCTC

Isolation

Human

References

References and other information relating to this product are available online at www.atcc.org.

Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate. This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials. Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at www.atcc.org

Additional information on this culture is available on the ATCC web site at www.atcc.org.
© ATCC 2016. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [11/10]

American Type Culture Collection
PO Box 1549
Manassas, VA 20108 USA
www.atcc.org

800.638.6597 or 703.365.2700
Fax: 703.365.2750
Email: Tech@atcc.org

Or contact your local distributor