



## Product Sheet

# MEF (CF-1) (ATCC® SCRC-1040™)

### Please read this FIRST



Storage Temp.  
**liquid nitrogen**  
vapor phase

---



Biosafety Level  
**1**

### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium:

- fetal bovine serum to a final concentration of 15%

This medium is formulated for use with a 5% CO<sub>2</sub> in air atmosphere. (Standard DMEM formulations contain 3.7 g/L sodium bicarbonate and a 10% CO<sub>2</sub> in air atmosphere is then recommended).

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: MEF (CF-1) (ATCC® SCRC-1040™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

## Description

**Organism:** *Mus musculus*, mouse

**Strain:** CF-1, non-inbred mouse strain (non-agouti albino) from Carworth Farms

**Tissue:** Embryo

**Cell Type:** Fibroblast

**Age:** 14 days gestation embryo

**Gender:** male and female mixed

**Morphology:** Fibroblast

**Growth Properties:** Adherent

## Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

## SAFETY PRECAUTION

ATCC highly recommends that protective gloves and clothing always be used and a full face mask always be worn when handling frozen vials. It is important to note that some vials leak when submersed in liquid nitrogen and will slowly fill with liquid nitrogen. Upon thawing, the conversion of the liquid nitrogen back to its gas phase may result in the vessel exploding or blowing off its cap with dangerous force creating flying debris.

## Unpacking & Storage Instructions

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below -130°C, preferably in liquid nitrogen vapor, until ready for use.

## Handling Procedure for Frozen Cells

To insure the highest level of viability, be sure to warm media to 37°C before using it on the cells.

Flasks do not need to be coated before plating MEFs.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water.
2. Remove the vial from the water bath as soon as the contents are half way thawed (approximately 90 seconds), and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
3. Transfer the vial's contents plus 5 mL of complete DMEM to a 15 mL centrifuge tube. Use an additional 1 mL of media to rinse the vial and transfer the liquid to the 15 mL tube. Add 4 mL of complete DMEM to bring the total volume to 10 mL.
4. Gently mix and pellet the cells by centrifugation @ 270 x g for 5 minutes.
5. Discard the supernatant and resuspend the cells with 10 mL fresh growth medium (warm) and plate the cells at seed density of 0.8 X 10<sup>4</sup> cells/cm<sup>2</sup>.
6. Add fresh growth medium (warm) to the appropriate size flask.
7. Incubate 37°C in a 5% CO<sub>2</sub> in air atmosphere.
8. Fluid change twice a week or when pH decreases.

*It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).*

## Subculturing Procedure

To insure the highest level of viability, be sure to warm media and Trypsin / EDTA to 37°C before using it on the cells. Cells should be split when they reach confluency. Split cells at approximately 0.4 X 10<sup>4</sup> cells/cm<sup>2</sup>


1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 1XPBS (SCRR-2201) solution to remove all traces of serum, which contain trypsin inhibitor.
3. Add 5 mL of Trypsin-EDTA (0.25% (w/v) Trypsin-0.53 mM EDTA solution, ATCC# 30-2101) solution to flask and incubate for 1 minute, gently tapping the flask observe cells under an inverted microscope until cells detach (usually within 1 to 2 minutes).
4. Add 6.0 to 8.0 mL of complete growth medium and rinse surface of the flask to detach all cells. Gently pipetting up and down will break cell clumps.
5. Transfer all cells into a centrifuge bottle or tube and centrifuge at 271x g for 5 minutes.
6. Remove and discard the supernatant




Product Sheet

# MEF (CF-1) (ATCC® SCRC-1040™)

Please read this FIRST

 Storage Temp.  
**liquid nitrogen vapor phase**

---

 Biosafety Level  
**1**

### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Complete Growth Medium

The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium:

- fetal bovine serum to a final concentration of 15%

This medium is formulated for use with a 5% CO<sub>2</sub> in air atmosphere. (Standard DMEM formulations contain 3.7 g/L sodium bicarbonate and a 10% CO<sub>2</sub> in air atmosphere is then recommended).

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: MEF (CF-1) (ATCC® SCRC-1040™)

7. Add 10 mL complete growth medium to cell pellet and with 10 mL pipette resuspend the cells gently (create a single-cell suspension).
8. Add more complete growth medium to cell suspension as needed to plate cells.
9. Place flasks in incubator @ 37°C with a 5% CO<sub>2</sub> in air atmosphere.

**Note:** For more information on enzymatic dissociation and subculturing of cell lines consult Chapter 13 in *Culture of Animal Cells, a manual of Basic Technique* by R. Ian Freshney, 5th edition, published by Alan R. Liss, N.Y., 2005.

Flask/Plate	Growth Area (cm <sup>2</sup> )	1xPBS (mL)	Trypsin/EDTA (mL)	Equal vol. Complete Growth Medium (mL)	Growth Medium (mL)
T225	225	10 ± 0.2	6 ± 0.2	6 ± 0.2	30
75	75	5 ± 0.1	3 ± 0.1	3 ± 0.1	12
T25	25	3 ± 0.1	1.5 ± 0.1	1.5 ± 0.1	6
6 well	9.5	1 ± 0.1	1 ± 0.1	1 ± 0.1	3

**Subcultivation Ratio:** Plate the cells at approximately of 0.4 X 10<sup>4</sup> cells/cm<sup>2</sup>.

**Medium Renewal:** Twice a week or when pH decreases.

### Comments

The growth of these cells should be arrested before being used as a feeder layer. ATCC has successfully irradiated (SCRC-1040.1™) and treated the cells with Mitomycin C (SCRC-1040.2a™) for use as a feeder layer. If the MEFs are being used as a feeder layer for ES cells, it is not recommended to use them past passage no. 6 (P6).

### References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

### Biosafety Level: 1

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

### ATCC Warranty

ATCC® products are warranted for 30 days from the date of shipment, and this warranty is valid only if the product is stored and handled according to the information included on this product information sheet. If the ATCC® product is a living cell or microorganism, ATCC lists the media formulation that has been found to be effective for this product. While other, unspecified media may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this product. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

### Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor



Product Sheet

**MEF (CF-1) (ATCC® SCRC-1040™)**

patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.


This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of materials on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of such materials.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).


© ATCC 2014. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [01/06]

**Please read this FIRST**



Storage Temp.  
**liquid nitrogen**  
**vapor phase**

---



Biosafety Level  
**1**

**Intended Use**

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

**Complete Growth Medium**

The base medium for this cell line is ATCC-formulated Dulbecco's Modified Eagle's Medium, Catalog No. 30-2002. To make the complete growth medium, add the following components to the base medium:

- fetal bovine serum to a final concentration of 15%

This medium is formulated for use with a 5% CO<sub>2</sub> in air atmosphere. (Standard DMEM formulations contain 3.7 g/L sodium bicarbonate and a 10% CO<sub>2</sub> in air atmosphere is then recommended).

**Citation of Strain**

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: MEF (CF-1) (ATCC® SCRC-1040™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

*Or contact your local distributor*