





Product Sheet

# Canine parvovirus (ATCC® VR-953™)

Please read this **FIRST**



---



## Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

## Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: Canine parvovirus (ATCC® VR-953™)

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor

## Description

**Strain:** C-780916

**Classification:** *Parvoviridae*, *Parvovirus*, Feline panleukopenia virus

**Original Source:** Feces of dog with hemorrhagic enteritis, 1978

**Depositor:** LE Carmichael, New York State College of Veterinary Medicine, Cornell University

## Batch-Specific Information

Refer to the Certificate of Analysis for batch-specific test results.

## Propagation

**Propagation Host:**

A-72 (ATCC® CRL-1542™)

**Effect on Host:**

CPE, refractile cells, cell rounding and sloughing

**Growth Conditions**

**Recommendations for Infection:** Plate cells 2-4 hours prior to infection and infect when cultures are 30-40% confluent. Remove medium and inoculate with a small volume of virus (e.g. 1 mL per 25 cm<sup>2</sup>) diluted to provide an optimal MOI (e.g. 0.1). Adsorb 1-2 hours at 35°C in a humidified atmosphere, rocking every 15 minutes to redistribute inoculum. End adsorption by adding virus growth medium.

**Incubation:** Incubate infected culture for 3-9 days at 35°C in a humidified atmosphere, until CPE are well advance throughout.

**Virus Growth Medium:** Leibovitz's L-15 Medium (ATCC 30-2008) + 2% FBS (ATCC 30-2020)

## Comments

Non-pathogenic for neonatal pups. Antigenically indistinguishable from Feline panleukopenia virus and Mink enteritis virus, but has broader host range and different HA spectrum, especially in regards to pH conditions. Hemagglutinates porcine, Rh+ monkey, and feline red blood cells at pH 6.0-7.5. May fail to immunize dogs if passaged 8-10 times in feline cells.

## References

References and other information relating to this product are available online at [www.atcc.org](http://www.atcc.org).

## Key Abbreviations

A/72, canine fibroma cell line

°C, degrees Celsius

CO<sub>2</sub> (CO2), carbon dioxide

CPE, cytopathic effect

CRFK, Crandell feline kidney cells

HA, hemagglutination

MOI, multiplicity of infection

Mv 1 Lu, mink lung epithelial cells

RBC, red blood cells

MOI, multiplicity of infection

## Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the current publication of the *Biosafety in Microbiological and Biomedical Laboratories* from the U.S. Department of Health and Human Services Centers for Disease Control and Prevention and National Institutes for Health.

## ATCC Warranty



The viability of ATCC® products is warranted for 30 days from the date of shipment, and is valid only if the product is stored and cultured according to the information included on this product information sheet. ATCC lists the media formulation that has been found to be effective for this strain. While other, unspecified media



Product Sheet

## Canine parvovirus (ATCC® VR-953™)

### Please read this FIRST

	Storage Temp. <b>-70°C or colder</b>
.....	
	Biosafety Level <b>2</b>

### Intended Use

This product is intended for research use only. It is not intended for any animal or human therapeutic or diagnostic use.

### Citation of Strain

If use of this culture results in a scientific publication, it should be cited in that manuscript in the following manner: Canine parvovirus (ATCC® VR-953™)

may also produce satisfactory results, a change in media or the absence of an additive from the ATCC recommended media may affect recovery, growth and/or function of this strain. If an alternative medium formulation is used, the ATCC warranty for viability is no longer valid.

### Disclaimers

This product is intended for laboratory research purposes only. It is not intended for use in humans. While ATCC uses reasonable efforts to include accurate and up-to-date information on this product sheet, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, and use. ATCC is not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to insure authenticity and reliability of strains on deposit, ATCC is not liable for damages arising from the misidentification or misrepresentation of cultures.

Please see the enclosed Material Transfer Agreement (MTA) for further details regarding the use of this product. The MTA is also available on our Web site at [www.atcc.org](http://www.atcc.org)

Additional information on this culture is available on the ATCC web site at [www.atcc.org](http://www.atcc.org).

© ATCC 2017. All rights reserved. ATCC is a registered trademark of the American Type Culture Collection. [12/15]

American Type Culture Collection  
PO Box 1549  
Manassas, VA 20108 USA  
[www.atcc.org](http://www.atcc.org)

800.638.6597 or 703.365.2700  
Fax: 703.365.2750  
Email: [Tech@atcc.org](mailto:Tech@atcc.org)

Or contact your local distributor